

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A21D 2/24	A1	(11) International Publication Number: WO 99/62343 (43) International Publication Date: 9 December 1999 (09.12.99)
(21) International Application Number: PCT/DK99/00277 (22) International Filing Date: 21 May 1999 (21.05.99) (30) Priority Data: PA 1998 00749 29 May 1998 (29.05.98) DK 60/088,096 5 June 1998 (05.06.98) US (71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Corporate Patents, Novo Alle, DK-2880 Bagsværd (DK). (72) Inventor; and (75) Inventor/Applicant (for US only): ILUM NIELSEN, Ruby [DK/DK]; Gedebakken 9, DK-3520 Farum (DK).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> <i>With an indication in relation to deposited biological material furnished under Rule 13bis separately from the description.</i>
(54) Title: METHODS FOR USING XYLOGLUCAN ENDOTRANSGLYCOSYLASE IN BAKING (57) Abstract The present invention relates to methods for preparing a dough, comprising incorporating into the dough a composition comprising an effective amount of an XET which improves one or more properties of the dough or a baked product obtained from the dough. The present invention also relates to methods for preparing a baked product. The present invention also relates to compositions comprising an effective amount of an XET for improving one or more properties of a dough and/or a baked product obtained from the dough. The present invention further relates to doughs or baked products and to pre-mixes for a dough.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LJ	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

METHODS FOR USING XYLOGLUCAN ENDOTRANSGLYCOSYLASE IN BAKING

Field of the Invention

5 The present invention relates to methods for preparing a dough and/or baked product with a xyloglucan endotransglycosylase.

Background Art

10 The strength of a dough is an important aspect of baking for both small-scale and large-scale applications. A strong dough has a greater tolerance of mixing time, proofing time, and mechanical vibrations during dough transport, whereas a weak dough is less tolerant to these treatments. A strong dough with
15 superior rheological and handling properties results from flour containing a strong gluten network. Flour with a low protein content or a poor gluten quality results in a weak dough.

Dough "conditioners" are well known in the baking industry.

The addition of conditioners to bread dough has resulted in
20 improved machinability of the dough and improved texture, volume, flavour and freshness (increased resistance to staling) of the bread. Nonspecific oxidants, such as iodates, peroxides, ascorbic acid, potassium bromate and azodicarbonamide have a gluten strengthening effect. It has been suggested that these
25 conditioners induce the formation of interprotein bonds which strengthen the gluten, and thereby the dough. However, the use of several of the currently available chemical oxidizing agents has been met with consumer resistance or is not permitted by regulatory agencies.

30 The use of enzymes as dough conditioners has been considered as an alternative to the chemical conditioners. A number of enzymes have been used recently as dough and/or bread improving agents, in particular, enzymes which act on components present in large amounts in the dough. Examples of such enzymes are
35 found within the groups of amylases, proteases, glucose oxidases, and (hemi)cellulases, including pentosanases.

Xyloglucan endotransglycosylase (XET) is an enzyme which catalyses an endo-transglycosylation of xyloglucan, a structural

polysaccharide of plant cell walls. The enzyme is believed to be present in all plants, and in particular, land plants. XET has been extracted from dicotyledons, monocotyledons, such as graminaceous monocotyledons and liliaceous monocotyledons, and also from a moss and a liverwort (Fry, S., et al., 1992.. Biochem. J. 282: 821-828).

XET may be obtained from a plant as described in WO 95/13384, WO 97/23683, or EP 562 836, or it may be obtained as described by Fry, et al., op cit.

It is the object of the present invention to improve the properties of dough and/or baked products by the use of an XET.

Summary of the Invention

The present invention relates to methods for preparing a dough and a baked product made from dough, respectively, comprising incorporating into the dough an effective amount of a xyloglucan endotransglycosylase (XET).

The present invention also relates to methods for preparing a baked product.

The present invention also relates to compositions comprising an effective amount of one or more XETs, for improving one or more properties of a dough and/or a baked product obtained from the dough, and a carrier and/or a baking ingredient.

The present invention also relates to doughs and to baked products, respectively.

The present invention further relates to pre-mixes for a dough comprising an effective amount of an XET for improving one or more properties of a dough and/or a baked product obtained from the dough, and a carrier and/or a baking ingredient.

Detailed Description of the Invention

The present invention relates to methods for preparing a dough or a baked product made from dough comprising

incorporating into the dough an effective amount of an XET which improves one or more properties of the dough or the baked product obtained from the dough relative to a dough or a baked product in which XET is not incorporated.

5 In the methods of the present invention, an XET is incorporated into the dough by adding the XET to the dough, to any single ingredient from which the dough is to be made, and/or to any mixture of dough ingredients from which the dough is to be made. In other words, the XET may be added in any
10 step of the dough preparation and may be added in one, two, or more steps.

The term "effective amount" is defined herein as an amount of an XET which is sufficient for providing a measurable effect on at least one property of interest of the dough and/or baked
15 product.

The term "improved property" is defined herein as any property of a dough and/or a product obtained from the dough, particularly a baked product, which is improved by the action of an XET relative to a dough or product in which an XET is not
20 incorporated. The improved property may include, but is not limited to, increased strength of the dough, increased elasticity of the dough, increased stability of the dough, reduced stickiness of the dough, improved extensibility of the dough, improved machinability of the dough, increased volume of
25 the baked product, improved crumb structure of the baked product, improved softness of the baked product, improved flavour of the baked product, and/or improved antistaling of the baked product.

The use of an XET may result in an increased strength, stability, and/or reduced stickiness of the dough, resulting in
30 improved machinability, as well as in an increased volume and improved crumb structure and softness of the baked product. The effect on the dough may be particularly advantageous when a poor quality flour is used. The improved machinability is of
35 particular importance in connection with dough which is to be processed industrially.

The improved property may be determined by comparison of a dough and/or a baked product prepared without addition of an

XET in accordance with the methods of the present invention. Techniques which can be used to determine improvements achieved by use of the methods of present invention are described below in the Examples. Organoleptic qualities may be evaluated using
5 procedures well established in the baking industry, and may include, for example, the use of a panel of trained taste-testers.

The term "increased strength of the dough" is defined herein as a the property of dough which has generally more elastic
10 properties and/or requires more work input to mould and shape.

The term "increased elasticity of the dough" is defined as the property of a dough which has a higher tendency to regain its original shape after being subjected to a certain physical strain.

15 The term "increased stability of the dough" is defined herein as the property of a dough that is less susceptible to mechanical abuse thus better maintaining its shape and volume.

The term "reduced stickiness of the dough" is defined herein as the property of a dough that has less tendency to adhere to
20 surfaces, e.g., in the dough production machinery, and is either evaluated empirically by the skilled test baker or measured by the use of a texture analyzer (e.g., TAXT2) as known in the art.

The term "improved extensibility of the dough" is defined as
25 the property of a dough that can be subjected to increased strain or stretching without rupture.

The term "improved machinability of the dough" is defined herein as the property of a dough that is generally less sticky and /or more firm and /or more elastic.

30 The term "increased volume of the baked product" is measured as the specific volume of a given loaf of bread (volume/weight) determined typically by the traditional rapeseed displacement method.

The term "improved crumb structure of the baked product" is
35 defined herein as the property of a baked product with finer and/or thinner cell walls in the crumb and/or more uniform/homogenous distribution of cells in the crumb and is usually evaluated empirically by the skilled test baker.

The term "improved softness of the baked product" is the opposite of "firmness" and is defined herein as the property of a baked product that is more easily compressed and is evaluated either empirically by the skilled test baker or measured by the use of a texture analyzer (e.g., TAXT2) as known in the art.

The term "improved flavour of the baked product" is evaluated as mentioned above by a trained test panel.

The term "improved antistaling of the baked product" is defined herein as the properties of a baked product that has a reduced rate of deterioration of quality parameters, e.g., softness and/or elasticity, during storage.

In a preferred embodiment, the XET improves one or more properties of the dough or the baked product obtained from the dough. In another preferred embodiment, the XET improves one or more properties of the dough and the baked product obtained from the dough.

In a preferred embodiment, the improved property is increased strength of the dough. In another preferred embodiment, the improved property is increased elasticity of the dough. In another preferred embodiment, the improved property is increased stability of the dough. In another preferred embodiment, the improved property is reduced stickiness of the dough. In another preferred embodiment, the improved property is improved extensibility of the dough. In another preferred embodiment, the improved property is improved machinability of the dough. In another preferred embodiment, the improved property is increased volume of the baked product.

In another preferred embodiment, the improved property is improved crumb structure of the baked product. In another preferred embodiment, the improved property is improved softness of the baked product. In another preferred embodiment, the improved property is improved flavour of the baked product.

In another preferred embodiment, the improved property is improved antistaling of the baked product.

The term "dough" is defined herein as a mixture of flour and other ingredients firm enough to knead or roll. The dough may be fresh, frozen, par-baked, or pre-baked. Preferably, the dough of the present invention is fresh or par-baked, i.e., the

dough is preferably baked without being frozen. The preparation of frozen dough is described by Kulp and Lorenz in *Frozen and Refrigerated Doughs and Batters*.

The term "baked product" is defined herein as any product prepared from dough, either of a soft or a crisp character. Examples of baked products, whether of a white, light or dark type, which may advantageously be produced by the present invention are bread (in particular white, whole-meal or rye bread), typically in the form of loaves or rolls, French baguette-type bread, noodles, pasta, pizzas, pita bread, tortillas, tacos, cakes, pancakes, biscuits, cookies, pie crusts, steamed bread, and crisp bread, and the like.

The XET may be any XET which provides an improved property to a dough and/or to a baked product obtained from the dough.

The term "xyloglucan endotransglycosylase (XET)" as used in the present invention is defined herein as glycosyltransferase which has the capacity to transfer a high molecular weight portion from a donor xyloglucan to a suitable acceptor such as a xyloglucan-derived monosaccharide.

Examples of an XET useful in the methods of the present inventions are described in PCT/DK98/00076, WO 95/13384 and WO 97/23683.

In the methods of the present invention, any XET may be used which possesses suitable enzyme activity in a pH and temperature range appropriate for making a dough and/or a baked product. It is preferable that the XET is active over broad pH and temperature ranges.

In a preferred embodiment, the XET has a pH optimum in the range of about 3 to about 10. In a more preferred embodiment, the XET has a pH optimum in the range of about 4.5 to about 8.5.

In a preferred embodiment, the XET has a temperature optimum in the range of about 5°C to about 100°C. In a more preferred embodiment, the XET has a temperature optimum in the range of about 25°C to about 75°C.

The source of the XET to be used according to the present invention is not critical for improving one or more properties of a dough and/or a baked product. Accordingly, the XET may be

obtained from any source such as a plant, microorganism, or animal. The XET is preferably obtained, e.g., from a microbial source, such as a bacterium or a fungus, e.g., a filamentous fungus or a yeast.

- 5 In a preferred embodiment, the XET is obtained from a plant source. For example, the XET may be obtained from a dicotyledon or a monocotyledon, in particular a dicotyledon selected from the group consisting of cauliflowers, soy beans, tomatoes, potatoes, rapeseeds, sunflowers, cotton, and tobacco, or a
10 monocotyledon selected from the group consisting of wheat, rice, corn and sugar cane.

In another preferred embodiment, the XET is obtained from a fungal source, and more preferably from a fungus from a Basidiomycota, Zygomycota, Ascomycota or Mitosporic strain.

- 15 A preferred Basidiomycota strain is a Hymenomycetes strain belonging to the orders Coriolales, Schizophyllales, Stereales or Xenasmatales; in particular a strain belonging to one of the families Coriolaceae, Corticiaceae, Schizophyllaceae, Stereaceae or Tubulicrinaceae. Preferred genera is one of the following:
20 *Trametes*, *Corticium*, *Schizophyllum*, or *Tubulicrinis*. A preferred species is one of the following: *Trametes hirsuta*, *Corticium roseum*, *Schizophyllum* sp, *Stereum hirsutum* or *Tubulicrinis subulatus*.

- Preferred Ascomycota are strains belonging to the classes
25 *Loculoascomycetes*, *Discomycetes*, *Pyrenomycetes*, and *Plectomycetes*, preferably those belonging to the orders *Dothideales*, *Rhytismatales*, *Pezizales*, *Leotiales*, *Xylariales*, *Hypocreales*, *Halosphaeriales*, *Phyllachorales*, *Diaporthales* and *Eurotiales*.

- Preferred strains are strains belonging to the families
30 *Botryosphaeriaceae*, *Dothioraceae*, *Mycosphaerellaceae*, *Tubeufiaceae*, *Pleosporaceae*, *Leptosphaeriaceae*, *Rhytismataceae*, *Sarcosomataceae*, *Pyronemataceae*, *Ascobolaceae*, *Sclerotiniaceae*, *Amphisphaeriaceae*, *Xylariaceae*, *Hypocreaceae*, *Halosphaeriaceae*, *Phyllachoraceae*, *Valsaceae*, *Melanconidaceae* and *Trichocomataceae*; especially strains belonging to the genera *Diplodia*,
35 *Plowrightia*, *Phyllosticta*, *Septoria*, *Tubeufia*, *Alternaria*, *Coniothyrium*, *Phoma*, *Embellisia*, *Tiarosporella*, *Galiella*, *Pseudoplectania*, *Pyronema*, *Oedocephalum*, *Botrytis*, *Aposphaeria*,

Pestalotia, Pestalotiopsis, Poronia, Nodulisporium, Xylaria, Fusarium, Verticillium, Volutella, Chaetapio-
spora, Lulworthia, Colletotrichum, Cytospora, Discula, Phomopsis, Coryneum, Seimatosporium, Aspergillus, Eurotium, Eupenicillium, Peni-
cillium, Petromyces and Talaromyces.

In another preferred embodiment the XET is obtained from a Diplodia gossypina, Plowrightia ribesia, Phyllosticta sp, Septoria sp, Tubeufia amazonensis, Alternaria sp, Embellisia hyacinthi, Phoma neoloba, Phoma tropica, Coniothyrium sp, Coniothyrium olivaceum, Coniothyrium dunckii, Galiella celebica, Pseudoplectania nigrella, Pyronema domesticum, Oedocephalum sp, Botrytis cinerea, Aposphaeria sp, Pestalotia sp, Pestalotiopsis sp. Poronia punctata, Xylaria sp, Nodulisporium sp, Fusarium solani, Verticillium sp, Volutella buxi, Chaetapio-
spora rhododendri, Lulworthia uniseptata, Colletotrichum aculatum, Colletotrichum crassipes, Cytospora spp, Discula sp, Phomopsis ilicis, Phomopsis cirsii, Coryneum castaneicola, Seimatosporium lichenicola, Aspergillus tamaris, Eurotium chevalieri, Eupenicillium javanicum, Penicillium capsulatum, Penicillium olsonii, Penicillium pinophilum, Penicillium roqueforti, Penicillium italicum, Penicillium canescens, Penicillium verruculosum, Petromyces alliaceus and Talaromyces flavus.

In another preferred embodiment the XET is obtained from a strain of Tiarosporella. In an even more preferred embodiment the XET is obtained from a strain of Tiarosporella phaseolina, or a synonym or teleomorph thereof. A strain of Tiarosporella phaseolina (Macrophomina sp., Classification: Ascomycota, Discomycetes, Rhytismatales, Rhytismataceae) has been deposited according to the Budapest Treaty on the International Recognition of the Deposits of Microorganisms for the Purpose of Patent Procedures, on 28 January 1997, at Centraalbureau voor Schimmelcultures (CBS) under Accession No. CBS 446.97. In a most preferred embodiment the XET is obtained from Tiarosporella phaseolina CBS No. 446.97.

Examples of useful Zygomycota are strains belonging to the order Mucorales, preferably strains belonging to the families Chaetocladiaceae and Mucoraceae.

Preferred strains belong to the genera *Dichotomocladium*, *Actinomucor*, *Gongronella*, *Sporodiniella* and *Mucor*, in particular *Dichotomocladium hesseltinei*, *Actinomucor elegans*, *Gongronellla butleri*, *Sporodiniella umbellata* and *Mucor miehei* var *minor*.

An example of a strain of uncertain taxonomy from which an XET may be obtained is *Vialaea insculpta*.

Examples of strains belonging to the Mitosporic fungi are *Acrodontium crateriforme*, *Aureobasidium pullulans*, *Circinotrichum* sp, *Cryptocline* sp, *Ellisiopsis* sp, *Epicoccum nigrum*, *Gliocladium* sp, *Helicorhoidion irregulare*, *Hendersonia* spp, *Mariannaea* sp, *Microsphaeropsis* sp, *Ramularia* sp, *Sarcopodium* sp, *Spadicoides sto*, *Speiropsis pedatospora*, *Sporotrichum exile*, *Stilbella* sp, *Trichothecium* sp, *Trimmatostroma abietes*, *Tubakia dryina*, *Wiesneriomyces* sp and *Zygosporium masonii*.

In a preferred embodiment, the XET is obtained from a bacterial source. For example, the XET may be obtained from a gram-negative or gram-positive bacterial cell.

Examples of XET-producing gram-positive bacteria are strains belonging to the genus *Bacillus*.

In another preferred embodiment, the XET may be obtained from the following strains which have been found to be XET-positive:

1. *Dichotomocladium hesseltinei*. Acc No of strain: CBS 164.61.
Classification: Zygomycota, Mucorales, Chaetocladiaceae.
2. *Actinomucor elegans*. Ex of Acc No of strain: CBS 154.86.
Classification: Zygomycota, Mucorales, Chaetocladiaceae.
3. *Mucor miehei* var *minor*. Acc No of strain: ATCC 36018.
Classification: Zygomycota, Mucorales, Mucoraceae.
4. *Gongronella butleri*. A strain of *Gongronella butleri* has been deposited according to the Budapest Treaty on the International Recognition of the Deposits of Microorganisms for the Purpose of Patent Procedures, on 28 January 1997, at Centraalbureau voor Schimmelcultures (CBS), under Accession No. CBS 448.97. Classification: Zygomycota, Mucorales,

Chaetocladiaceae.

5. *Sporodiniella umbellata*. Acc No of strain: CBS 195.77.

Classification: Zygomycota, Mucorales, Mucoraceae.

6. *Phyllosticta* sp. Isolated from a leaf of *Pithecolobium* sp.,

5 China. Classification: Ascomycota, Loculoascomycetes, Dothidiales, Mycosphaellaceae.

7. *Septoria* sp. A strain of *Septoria* sp has been deposited according to the Budapest Treaty on the International

10 Recognition of the Deposits of Microorganisms for the Purpose of Patent Procedures, on 2 January 1996, at Centraalbureau voor Schimmelcultures (CBS), under Accession No. CBS 831.95. Classification: Ascomycota, Loculoascomycetes, Dothidiales, Mycosphaellaceae.

8. *Diplodia gossypina*. A strain of *Diplodia gossypina* has been

15 deposited according to the Budapest Treaty on the International Recognition of the Deposits of Microorganisms for the Purpose of Patent Procedures; on 12 March 1996, at Centraalbureau voor Schimmelcultures (CBS), under Accession No. CBS 274.96. Classification: Ascomycota, Loculoascomycetes, Dothidiales,

20 Botryosphaeriaceae.

9. *Plowrightia ribesia*. Isolated from *Ribes* sp., Denmark.

Classification: Ascomycota, Loculoascomycetes, Dothidiales, Dothioraceae.

10. *Tubeufia amazonensis*. Acc No of Strain: ATCC 42524.

25 Classification: Ascomycota, Loculoascomycetes, Dothidiales, Tubeufiaceae.

11. *Alternaria* sp. Classification: Ascomycota, Loculoascomycetes, Dothidiales, Pleosporaceae.

12. *Embellisia hyacinthi*. Acc No of species IMI 211561.

30 Classification: Ascomycota, Loculoascomycetes, Dothidiales, Pleosporaceae.

13. *Phoma neoloba*. Classification: Ascomycota, Loculoascomycetes, Dothideales, Pleosporaceae.
14. *Phoma tropica*. Ex on Acc No of species CBS 537.66. Classification: Ascomycota, Loculoascomycetes, Dothideales, Pleosporaceae.
15. *Coniothyrium* sp. Classification: Ascomycota, Loculoascomycetes, Dothideales, Leptosphaeriaceae.
16. *Coniothyrium olivaceum*. Ex on Acc No of species CBS 304.68. Classification: Ascomycota, Loculoascomycetes, Dothideales, Leptosphaeriaceae.
17. *Coniothyrium dunckii*. Classification: Ascomycota, Loculoascomycetes, Dothideales, Leptosphaeriaceae.
18. *Tiarosporella* sp. Classification: Ascomycota, Discomycetes, Rhytismatales, Rhytismataceae.
19. *Galiella celebica*. Isolated from a sample collected in Japan. Classification: Ascomycota, Discomycetes, Pezizales, Sarcosomataceae.
20. *Pseudoplectania nigrella*. A strain of *Pseudoplectania nigrella* has been deposited according to the Budapest Treaty on the International Recognition of the Deposits of Microorganisms for the Purpose of Patent Procedures, on 28 January 1997, at Centraalbureau voor Schimmelcultures (CBS), under Accession No. CBS 444.97. Classification: Ascomycota, Discomycetes, Pezizales, Sarcosomataceae.
21. *Pyronema domesticum*. Isolated from a sample collected in Norway. Classification: Ascomycota, Discomycetes, Pezizales, Pyronemataceae.
22. *Oedocephalum* sp. Classification: Ascomycota, Discomycetes, Pezizales, Ascobolaceae.
23. *Botrytis cinerea*. A strain of *Botrytis cinerea* has been deposited according to the Budapest Treaty on the

International Recognition of the Deposits of Microorganisms for the Purpose of Patent Procedures, on 28 January 1997, at Centraalbureau voor Schimmelcultures (CBS), under Accession No. CBS 447.97. Classification: Ascomycota, Discomycetes, 5 Leotiales, Sclerotiniaceae.

24. *Aposphaeria* sp. Classification: Ascomycota, Discomycetes, Leotiales, Sclerotiniaceae.

25. *Pestalotia* sp. A strain of *Pestalotia* sp. has been deposited according to the Budapest Treaty on the International 10 Recognition of the Deposits of Microorganisms for the Purpose of Patent Procedures, on 28 January 1997, at Centraalbureau voor Schimmelcultures (CBS), under Accession No. CBS 445.97. Classification: Ascomycota, Pyrenomycetes, Xylariales, Amphisphaeriaceae.

15 26. *Pestalotiopsis* sp. Classification: Ascomycota, Pyrenomycetes, Xylariales, Amphisphaeriaceae.

27. *Poronia punctata*. Isolated from a sample collected in Sweden. Classification: Ascomycota, Pyrenomycetes, Xylariales, Xylariaceae.

20 28. *Xylaria* sp. Isolated from a leaf of the palm, *Sabal jamaicensis*, growing in Mona, Jamaica. Classification: Ascomycota, Pyrenomycetes, Xylariales, Xylariaceae Xylariaceae.

29. *Nodulisporium* sp. Classification: Ascomycota, Pyrenomycetes, Xylariales, Xylariaceae.

25 30. *Fusarium solani*. Isolated from a sample of grain of maize collected in India. Classification: Ascomycota, Pyrenomycetes, Hypocreales, Hypocreaceae.

31. *Verticillium* sp. A strain of *Verticillium* sp. has been deposited according to the Budapest Treaty on the International 30 Recognition of the Deposits of Microorganisms for the Purpose of Patent Procedures, on 2 January 1996, at Centraalbureau voor Schimmelcultures (CBS), under Accession No. CBS 830.95.

Classification: *Ascomycota*, *Pyrenomycetes*, *Hypocreales*,
Hypocreaceae.

32. *Volutella buxi*. Acc No of Strain: IMI 049467.

Classification: *Ascomycota*, *Pyrenomycetes*, *Hypocreales*,
5 *Hypocreaceae*.

33. *Chaetapiospora rhododendri*. Classification: *Asco-*
mycota, *Pyrenomycetes*, *Xylariales*, *Xylariaceae*,
Hyponectriaceae.

34. *Lulworthia uniseptata*. A strain of *Lulworthia*
10 *uniseptata* has been deposited according to the Budapest Treaty
on the International Recognition of the Deposits of Micro-
organisms for the Purpose of Patent Procedures, on 28 January
1997, at Centraalbureau voor Schimmelcultures (CBS), under
Accession No. CBS 442.97. Classification: *Ascomycota*,
15 *Pyrenomycetes*, *Halosphaeriales*, *Halosphaeriaceae*.

35. *Colletotrichum aculatum*. Classification: *Ascomycota*,
Pyrenomycetes, *Phyllachorales*, *Phyllachoraceae*.

36. *Colletotrichum crassipes*. Classification: *Ascomycota*,
Pyrenomycetes, *Phyllachorales*, *Phyllachoraceae*.

20 37. *Cytospora* sp. A strain of *Cytospora* sp has been
deposited according to the Budapest Treaty on the International
Recognition of the Deposits of Microorganisms for the Purpose
of Patent Procedures, on 23 January 1997, at Centraalbureau
voor Schimmelcultures (CBS), under Accession No. CBS 424.97.

25 Classification: *Ascomycota*, *Pyrenomycetes*, *Diaporthales*,
Valsaceae.

38. *Cytospora* sp. A strain of *Cytospora* sp has been
deposited according to the Budapest Treaty on the International
Recognition of the Deposits of Microorganisms for the Purpose
30 of Patent Procedures, on 23 January 1997, at Centraalbureau
voor Schimmelcultures (CBS), under Accession No. CBS 425.97.
Classification: *Ascomycota*, *Pyrenomycetes*, *Diaporthales*,
Valsaceae.

39. *Discula* sp. Classification: Ascomycota, Pyrenomycetes, Diaporthales, Valsaceae.
40. *Phomopsis ilicis*. Classification: Ascomycota, Pyrenomycetes, Diaporthales, Valsaceae.
- 5 41. *Phomopsis cirsii*. Classification: Ascomycota, Pyrenomycetes, Diaporthales, Valsaceae.
42. *Coryneum castaneicola*. Classification: Ascomycota, Pyrenomycetes, Diaporthales, Melanconidaceae.
- 10 43. *Seimatosporium lichenicola*. Classification: Ascomycota, Pyrenomycetes, Diaporthales, Melanconidaceae.
44. *Aspergillus tamarii*. Ex of Acc No of strain: CBS 821.72. Classification: Ascomycota, Plectomycetes, Eurotiales, Trichocomataceae.
- 15 45. *Eurotium chevalieri*. Ex of Acc No of strain: CBS 472.91. Classification: Ascomycota, Plectomycetes, Eurotiales, Trichocomataceae.
46. *Penicillium capsulatum*. Ex of Acc No of strain: CBS 273.86. Classification: Ascomycota, Plectomycetes, Eurotiales, Trichocomataceae.
- 20 47. *Penicillium olsonii*. Ex of Acc No of strain: CBS 523.89. Classification: Ascomycota, Plectomycetes, Eurotiales, Trichocomataceae.
48. *Penicillium pinophilum*. Ex of Acc No of strain: CBS 440.89. Classification: Ascomycota, Plectomycetes, Eurotiales, 25 Trichocomataceae.
49. *Penicillium roqueforti*. Ex of Acc No of strain: CBS 167.91. Classification: Ascomycota, Plectomycetes, Eurotiales, Trichocomataceae.
50. *Penicillium italicum*. Ex of Acc No of strain: IMI 078 30 681. Classification: Ascomycota, Plectomycetes, Eurotiales,

Trichocomataceae.

51. *Penicillium canescens*. Ex of Acc No of strain: CBS 579.70. Isolated from a salt mine in Egypt. Classification: *Ascomycota, Plectomycetes, Eurotiales, Trichocomataceae.*
- 5 52. *Eupenicillium javanicum*. Ex of Acc No of the strain: CBS 448.74. Classification: *Ascomycota, Plectomycetes, Eurotiales, Trichocomataceae.*
53. *Penicillium verruculosum*. Ex of Acc No of strain: CBS 563.92. Classification: *Ascomycota, Plectomycetes, Eurotiales,*
- 10 *Trichocomataceae.*
54. *Talaromyces flavus*. Acc No of the strain: ATCC 52201. Classification: *Ascomycota, Plectomycetes, Eurotiales, Trichocomataceae.*
55. *Petromyces alliaceus*. Acc No of strain: CBS 511.69.
- 15 Classification: *Ascomycota, Plectomycetes, Eurotiales, Trichocomataceae.*
56. *Trametes hirsuta*. Isolated from a sample collected in Denmark. Classification: *Basidiomycota, Hymenomycetes, Coriolales, Coriolaceae.*
- 20 57. *Schizophyllum* sp. A strain of *Schizophyllum* sp has been deposited according to the Budapest Treaty on the International Recognition of the Deposits of Microorganisms for the Purpose of Patent Procedures, on 28 January 1997, at Centraalbureau voor Schimmelcultures (CBS), under Accession No. CBS 443.97.
- 25 Classification: *Basidiomycota, Hymenomycetes, Schizophyllales, Schizophyllaceae.*
58. *Corticium roseum*. Isolated from a sample collected in Denmark. Classification: *Basidiomycota, Hymenomycetes, Aleurodiscales Cortiaceae.*
- 30 59. *Tubulicrinis subulatus*. Isolated from a sample collected in Denmark. Classification: *Basidiomycota, Hymenomycetes, Xenasmatales, Tubulicrinaceae.*

60. *Stereum hirsutum*. Isolated from a sample collected in Denmark. Classification: Basidiomycota, Hymenomycetes, Stereales, Stereaceae.

61. Strains of the Classification: Mitosporic fungus:
5 *Acrodontium crateriforme*, *Aureobasidium pullulans*,
Circinotrichum sp., *Cryptocline* sp., *Ellisiopsis* sp., *Epicoccum*
nigrum, *Gliocladium* sp., *Helicorhoidium irregulare*, *Hendersonia*
sp., *Mariannaea* sp., *Microsphaeropsis* sp., *Ramularia* sp.,
10 *Sarcopodium* sp., *Spadicoides* sp. Acc No of strain IMI203428,
Speiropsis pedatospora, *Sporotrichum exile*. Acc No of strain CBS
350.47, *Stilbella* sp., *Trichothecium* sp., *Trimmatostroma*
abietes, *Tubakia dryina*, *Wiesneriomyces* sp., and *Zygosporium*
masonii.

62. *Vialaea insculpta*. Classification: Uncertain.

15

63. *Bacillus alcalophilus*. A strain of *Bacillus alcalophilus*
has been deposited according to the Budapest Treaty on the
International Recognition of the Deposits of Microorganisms for
the Purpose of Patent Procedures, on 12 February 1997, at
20 Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH,
under Accession No. DSM 11404.

The taxonomic classification used herein builds primarily on
the system used in the NIH Database (Entrez, version spring
25 1996) available on World Wide Web at
<http://www.ncbi.nlm.nih.gov/Taxonomy/tax.html>.

Classification of organisms not included in the Entrez
database may be found in the following reference books which
are generally available and accepted in the art:

30 Ascomycetes: Eriksson, O.E. & Hawksworth, D.L.: *Systema*
Ascomycetum vol 12 (1993);

Basidiomycetes: Jülich, W.: *Higher Taxa of Basidiomycetes*,
Bibliotheca Mycologia 85, 485pp (1981);

Zygomycetes: O'Donnell, K.: *Zygomycetes in culture*,
35 *University of Georgia, US*, 257pp (1979).

General mycological reference books include: Hawksworth, D.L.,

Kirk, P.M., Sutton, B.C. and Pegler, D.N. *Dictionary of the fungi*, International Mycological Institute, 616 pp (1995); and Von Arx, J. A. *The genera of fungi sporulating in culture*, 424 pp (1981)..

5 The XET may be obtained from the organism in question by any suitable technique, and in particular by use of recombinant DNA techniques known in the art (c.f. Sambrook, J. et al., 1989, *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, NY, USA). The use of recombinant
10 DNA techniques generally comprises cultivation of a host cell transformed with a recombinant DNA vector, consisting of the product gene of interest inserted between an appropriate promoter and terminator, in a culture medium under conditions permitting the expression of the enzyme and recovering the
15 enzyme from the culture. The DNA sequence may be of genomic, cDNA or synthetic origin or any mixture of these, and may be isolated or synthesised in accordance with methods known in the art. The enzyme may also be obtained from its naturally occurring source, such as a plant or organism, or relevant part
20 thereof.

When an XET is added to dough intended for use in the preparation of baked products, it may exert a strengthening effect on dough constituents. The XET is used in an amount sufficient to provide the desired effect, i.e., the improved
25 properties in question. Thus, the dosage of the XET to be used in the methods of the present invention should be adapted to the nature and composition of the dough in question as well as to the nature of the XET to be used.

The term "composition" is defined herein as a dough-
30 improving and/or baked product-improving composition which, in addition to the XET, comprises one or more additional substances conventionally used in baking. The additional substance(s) may be other enzymes or chemical additives known in the art to be useful in dough preparation and/or baking.

35 The bread-improving and/or dough improving composition of the invention is generally included in the dough in an amount corresponding to 0.01-5%, in particular 0.1-3%. The XET is typically added in an amount corresponding to 0.01-100 mg

enzyme protein per kg of flour, preferably 0.1-25 mg enzyme protein per kg of flour, more preferably 0.1-10 mg enzyme protein per kg of flour, and most preferably 0.5-5 mg enzyme protein per kg of flour, particularly 1-5 mg/kg.

5 In terms of enzyme activity, the appropriate dosage of a given XET for exerting a desirable improvement of dough and/or baked products will depend on the enzyme and the enzyme substrate in question. The skilled person may determine a suitable enzyme unit dosage on the basis of methods known in
10 the art.

In a preferred embodiment, the dough-improving and/or baked product-improving composition of the invention comprises an XET selected from the group consisting of:

(a) a polypeptide which is encoded by a nucleic acid sequence
15 which hybridizes under low stringency conditions with (i) the nucleic acid sequence set forth in SEQ ID NO:1, (ii) its complementary strand, or (iii) a subsequence thereof;

(b) a polypeptide having an amino acid sequence which has at least 50% identity with the amino acid sequence set forth in
20 SEQ ID NO:2;

(c) an allelic variant of (a) or (b); and

(d) a fragment of (a), (b) or (c), wherein the fragment retains XET activity.

25 Hybridisation indicates that by methods of standard Southern blotting procedures (for example, as described by J. Sambrook, E.F. Fritsch, and T. Maniatus (1989, *Molecular Cloning, A Laboratory Manual*, 2d edition, Cold Spring Harbor, New York), the nucleic acid sequence hybridizes to an oligonucleotide
30 probe corresponding to the polypeptide encoding part of the nucleic acid sequence shown in SEQ ID NO:1, under low to high stringency conditions (i.e., prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 µg/ml sheared and denatured salmon sperm DNA, and either 25, 35 or 50% formamide for low,
35 medium and high stringencies, respectively). In order to identify a clone or DNA which is homologous with SEQ ID NO:1, the hybridisation reaction is washed three times for 30 minutes

each using 2 x SSC, 0.2% SDS preferably at least 50°C, more preferably at least 55°C, more preferably at least 60°C, more preferably at least 65°C, even more preferably at least 70°C, and most preferably at least 75°C.

5 The aforementioned probe can be considerably shorter than the entire sequence, but should be at least 15, preferably at least 25, and more preferably at least 40 nucleotides in length. Longer probes can also be used. Both DNA and RNA probes can be used. The probes are typically labelled for detecting
10 the corresponding gene (for example, with ³²P, ³H, ³⁵S, biotin, or avidin). For example, molecules to which a ³²P-, ³H- or ³⁵S-labelled oligonucleotide probe hybridizes may be detected by use of X-ray film.

The aforementioned polypeptide has an amino acid sequence
15 which has a degree of identity to the amino acid sequence of SEQ ID NO:2 of at least about 50%, preferably at least about 60%, preferably at least about 70%, more preferably at least about 80%, even more preferably at least about 90%, most preferably at least about 95%, and even most preferably at
20 least about 97%, which retain XET activity (hereinafter "homologous polypeptides"). In a preferred embodiment, the homologous polypeptides have an amino acid sequence which differs by five amino acids, preferably by four amino acids, more preferably by three amino acids, even more preferably by
25 two amino acids, and most preferably by one amino acid from the amino acid sequence of SEQ ID NO:2. For purposes of the present invention, the degree of identity between two amino acid sequences may be suitably determined by means of computer programs known in the art, such as GAP provided in the GCG
30 program package (Program Manual for the Wisconsin Package, Version 8, August 1994, Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA 53711) (Needleman, S.B. and Wunsch, C.D., (1970), Journal of Molecular Biology, 48, 443-453, using GAP with the following settings for polypeptide
35 sequence comparison: GAP creation penalty of 3.0 and GAP extension penalty of 0.1.

An allelic variant denotes any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic

variation arises naturally through mutation, and may result in phenotypic polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequences. The term
5 allelic variant is also used to denote a protein encoded by an allelic variant of a gene.

The XET and/or additional enzymes to be used in the methods of the present invention may be in any form suited for the use in question, e.g., in the form of a dry powder, agglomerated
10 powder, or granulate, in particular a non-dusting granulate, a liquid, in particular a stabilized liquid, or a protected enzyme. Granulates and agglomerated powders may be prepared by conventional methods, e.g., by spraying the XET onto a carrier in a fluid-bed granulator. The carrier may consist of
15 particulate cores having a suitable particle size. The carrier may be soluble or insoluble, e.g., a salt (such as NaCl or sodium sulfate), a sugar (such as sucrose or lactose), a sugar alcohol (such as sorbitol), starch, rice, corn grits, or soy. The XET and/or additional enzymes may be contained in slow-
20 release formulations. Methods for preparing slow-release formulations are well known in the art. Liquid enzyme preparations may, for instance, be stabilized by adding nutritionally acceptable stabilizers such as a sugar, a sugar alcohol or another polyol, lactic acid, or another organic acid
25 according to established methods. Granulated enzymes may be prepared according to the method disclosed in WO 97/423839.

For inclusion in pre-mixes or flour it is advantageous that the XET is in the form of a dry product, e.g., a non-dusting granulate, whereas for inclusion together with a liquid it is
30 advantageously in a liquid form.

A substrate of the XET in question may also be incorporated into the dough. The substrate may be incorporated into dough separately or together with the XET of interest, optionally as constituent(s) of the bread-improving and/or dough-improving
35 composition. Alternatively, an enzyme which acts on a substance endogenous to the flour to produce a substrate for the XET of interest may also be incorporated in the dough.

The specific amount of the substrate available for the XET of interest will depend on a number of factors, such as the baking process used, the length of time for mixing, fermentation, proofing and/or baking, the quality of the yeast and/or flour used, as well as the activity of endogenous and exogenous enzymes present.

A preferred substrate for XET is xyloglucan.

One or more additional enzymes may also be incorporated into the dough. The additional enzyme may be of any origin, including mammalian and plant, and preferably of microbial (bacterial, yeast or fungal) origin and may be obtained by techniques conventionally used in the art.

In a preferred embodiment, the additional enzyme may be an amylase, such as an alpha-amylase (useful for providing sugars fermentable by yeast and retarding staling), a beta-amylase, a cyclodextrin glucanotransferase, a peptidase, in particular, an exopeptidase (useful in flavour enhancement), a trans-lutaminase, a lipase (useful for the modification of lipids present in the dough or dough constituents so as to soften the dough), a phospholipase (useful for the modification of lipids present in the dough or dough constituents so as to soften the dough and improving gas retention in the dough), a cellulase, a hemicellulase, in particular a pentosanase such as xylanase (useful for the partial hydrolysis of pentosans which increases the extensibility of the dough), a protease (useful for gluten weakening in particular when using hard wheat flour), a protein disulfide isomerase, e.g., a protein disulfide isomerase as disclosed in WO 95/00636, a glycosyltransferase, and an oxidoreductase, e.g., a dehydrogenase, a cellobiose dehydrogenase, a fructose dehydrogenase or a lactate dehydrogenase; a peroxidase (useful for improving the dough consistency), a laccase, an aldose oxidase, a glucose oxidase, a pyranose oxidase, a lipoxygenase, an L-amino acid oxidase (useful in improving dough consistency) or a carbohydrate oxidase.

Commercially available amylases useful in the present invention are NOVAMYL™ (a *Bacillus stearothermophilus*

maltogenic amylase, available from Novo Nordisk A/S, Denmark), FUNGAMYL® (an *Aspergillus oryzae* alpha-amylase, available from Novo Nordisk A/S, Denmark), and BAN™ (a *Bacillus licheniformis* alpha-amylase, available from Novo Nordisk A/S, Denmark). A commercially available amyloglucosidase useful in the present invention is AMG™ (an *Aspergillus niger* amyloglucosidase, available from Novo Nordisk A/S, Denmark). Other useful commercially available amylase products include GRINDAMYL™ A 1000 or A 5000 (available from Grindsted Products, Denmark) and AMYLASE H or AMYLASE P (available from Gist-Brocades, The Netherlands). A commercially available glucose oxidase useful in the present invention is GLUZYME™ (an *Aspergillus niger* glucose oxidase, available from Novo Nordisk A/S, Denmark). Commercially available proteases useful in the present invention are NEUTRASE™ (a *Bacillus amyloliquefaciens* endoprotease, available from Novo Nordisk A/S, Denmark) and GLUTENASE™ (available from Novo Nordisk A/S, Denmark). A commercially available pentosanase useful in the present invention is PENTOPAN™ (a *Humicola insolens* pentosanase, available from Novo Nordisk A/S, Denmark). A commercially available lipase useful in the present invention is NOVOZYM® 677 BG (a *Thermomyces lanuginosus* lipase, available from Novo Nordisk A/S, Denmark).

When one or more additional enzyme activities are to be added in accordance with the methods of the present invention, these activities may be added separately or together with the XET, optionally as constituent(s) of the bread-improving and/or dough-improving composition. The other enzyme activities may be any of the above described enzymes and may be dosed in accordance with established baking practice.

In addition to the above mentioned additional enzymes, the XET may contain varying minor amounts of other enzymatic activities inherently produced by the producer organism in question.

In addition, or as an alternative, to additional enzyme components, one or more conventionally used dough and/or bread improving agents may also be incorporated into the dough. The agent may include proteins, such as milk powder (to provide

crust colour), gluten (to improve the gas retention power of weak flours), and soy (to provide additional nutrients and improve water binding); eggs such (either whole eggs, egg yolks or egg whites); fat such as granulated fat or shortening (to soften the dough and improve the texture of the bread); an emulsifier (to improve dough extensibility and, to some extent, the consistency of the resulting bread); an oxidant, e.g., ascorbic acid, potassium bromate, potassium iodate, azodicarbon amide (ADA) or ammonium persulfate (to strengthen the gluten structure); an amino acid, e.g., L-cysteine (to improve mixing properties); a sugar; a salt, e.g., sodium chloride, calcium acetate, sodium sulfate or calcium sulphate (to make the dough firmer); flour; and starch. Such components may also be added to the dough in accordance with the methods of the present invention.

Examples of suitable emulsifiers are mono- or diglycerides, diacetyl tartaric acid esters of mono- or diglycerides, sugar esters of fatty acids, polyglycerol esters of fatty acids, lactic acid esters of monoglycerides, acetic acid esters of monoglycerides, polyoxyethylene stearates, phospholipids, lysolecithin and lecithin.

The dough and/or baked product prepared by a method of the present invention may be based on wheat meal or flour, optionally in combination with other types of meal or flour such as corn meal, corn flour, rye meal, rye flour, oat meal, oat flour, soy meal, soy flour, sorghum meal, sorghum flour, potato meal, or potato flour.

The handling of the dough and/or baking may be performed in any suitable manner for the dough and/or baked product in question, typically including the steps of kneading the dough, subjecting the dough to one or more proofing treatments, and baking the product under suitable conditions, i.e., at a suitable temperature and for a sufficient period of time. For instance, the dough may be prepared by using a normal straight dough process, a sour dough process, an overnight dough method, a low-temperature and long-time fermentation method, a frozen dough method, the Chorleywood Bread process, or the Sponge and Dough process.

From the above disclosure it will be apparent that the dough of the invention is generally a leavened dough or a dough to be subjected to leavening. The dough may be leavened in various ways such as by adding sodium bicarbonate or the like, or by adding a leaven (fermenting dough), but it is preferable that the dough be leavened by adding a suitable yeast culture, such as a culture of *Saccharomyces cerevisiae* (baker's yeast). Any of the commercially available *Saccharomyces cerevisiae* strains may be employed.

The present invention also relates to the use of an XET for the preparation of pasta dough, preferably prepared from durum flour or a flour of comparable quality. The dough may be prepared by use of conventional techniques and the XET used in a similar dosage as that described above. The XET may be any of the types described above. When used in the preparation of pasta, the XET results in a strengthening of the gluten structure, a reduction in the dough stickiness, and an increased dough strength.

The present invention also relates to methods for preparing a baked product, comprising baking a dough produced by a method of the present invention to produce a baked product. The baking of the dough to produce a baked product may be performed using methods well known in the art.

The present invention also relates to dough and/or bread improving compositions comprising an effective amount of an XET for improving one or more properties of a dough and/or a baked product obtained from the dough, and a carrier and/or a baking ingredient. The compositions may further comprise a substrate for the XET, one or more additional enzymes, one or more conventionally used dough and/or bread improving agents, an enzyme which acts on a substance endogenous to the flour to produce a substrate for the XET, and/or a substance and the enzyme which acts on the substance to produce a substrate for the XET.

The present invention also relates to doughs and baked products, respectively, produced by the methods of the present invention.

The present invention further relates to a pre-mix, e.g., in the form of a flour composition, for dough and/or baked products made from dough, in which the pre-mix comprises an XET. The term "pre-mix" is defined herein to be understood in
5 its conventional meaning, i.e., as a mix of baking agents, generally including flour, which may be used not only in industrial bread-baking plants/facilities, but also in retail bakeries. The pre-mix may be prepared by mixing an XET or a bread-improving and/or dough-improving composition of the
10 invention comprising an XET with a suitable carrier such as flour, starch, a sugar or a salt. The pre-mix may contain other dough-improving and/or bread-improving additives, e.g., any of the agents, including enzymes, mentioned above.

The present invention further relates to baking additives in
15 the form of a granulate or agglomerated powder, which comprises the XET. The baking additive preferably has a narrow particle size distribution with more than 95% (by weight) of the particles in the range from 25 to 500 μm .

20 The present invention is further described by the following examples which should not be construed as limiting the scope of the invention.

EXAMPLES

5 Materials and Methods:

Preparation of White Bread (I)

10 The straight-dough bread-making method may be used according to AACC Method 10-10B (in Approved Methods of the American Association of Cereal Chemists, Ninth Edition, March 1995; AACC, St. Paul MN, USA).

Basic recipe

15	Wheat flour	100%
	Salt	1.5%
	Yeast (fresh)	5.3%
	Sugar	6.0%
20	Shortening	3.0%
	Water determined empirically	

25 All percentages are by weight relative to the wheat flour.

Procedure

1. Dough mixing (Hobart mixer):

The mixing time and speed should be determined by the skilled baker so as to obtain an optimum dough consistency
30 under the testing conditions used.

2. 1st punch (e.g., 52 minutes after start)

3. 2nd punch (e.g., 25 minutes later)

4. Moulding and panning (e.g., 13 minutes later).

5. Proofing to desired height (e.g., 33 minutes at 32°C, 82%
35 RH)

5. Baking (e.g., at 215°C for 24 minutes)

Preparation of White Bread (II)

40 The sponge-dough bread-making method may be used according to AACC Method 10-11 (in Approved Methods of the American Association of Cereal Chemists, Ninth Edition, March 1995;

AACC, St. Paul MN, USA).

Basic recipe for Sponge

	Wheat flour	60%
5	Yeast (compressed)	36%
	Yeast Food	2%
	Water	36%

All percentages are by weight relative to the wheat flour.

Procedure

1. Add water to compressed yeast
2. Add yeast food in dry form with flour
3. Mix sponge (Hobart A-120; Hobart Corp., Troy OH, USA):

15 0.5 minute at 1st speed

1 minute at 2nd speed

The mixing time may be adjusted so as to obtain an optimum dough consistency under the testing conditions used.

4. Ferment in a fermentation cabinet: 4 hours at 30°C, 85% RH

Basic recipe for Dough

	Wheat flour	40%
	Water	24%
	Sugar	5%
25	Shortening	3%
	Salt	2%

All percentages are by weight relative to the wheat flour.

Procedure

1. Add dough ingredients; begin mixer (1st speed)
2. Add sponge in three approximately equal portions at 15, 25, and 35 seconds mixing time; total mixing time: 1 minute
3. At 2nd speed, mix to obtain an optimum dough consistency
- 35 4. Ferment in a fermentation cabinet: 30 minutes at 30°C, 85% RH
5. Intermediate proof: 12-15 minutes in fermentation cabinet
6. Mould and final proof at 35.5°C, 92% relative humidity
7. Bake: 25 minutes at 218°C

Evaluation of Staling Properties of Bread

The degree of staling is determined on bread, e.g., on day

1, 3, 7 and 9 after baking. Evaluation of staleness and texture can be done according to AACC method 74-09. The principles for determination of softness and elasticity of bread crumb are as follows:

- 5 1. A slice of bread is compressed with a constant speed in a texture analyser, measuring the force for compression in g.
2. The softness of the crumb is measured as the force at 25% compression.
3. The force at 40% compression (P2) and after keeping 40%
10 compression constant for 30 seconds (P3) is measured. The ratio (P3/P2) is the elasticity of the crumb.

Preparation of White Layer Cake

The method may be used according to AACC Method 10-90 (in
15 Approved Methods of the American Association of Cereal Chemists, Ninth Edition, March 1995; AACC, St. Paul MN, USA).

Basic recipe

20	Flour	100%
	Sugar	140%
	Shortening	50%
	Non-fat Dry Milk	12%
	Dried Egg Whites	9%
25	Salt	3%
	Baking Powder, Water determined empirically.	

All percentages are by weight relative to the flour.

30 Procedure

1. Combine all dry ingredients and sift well
2. Add shortening and 60% of water
3. Mix at low speed for 0.5 minute in Hobart C-100 mixer
4. Mix at medium speed for 4 minutes
- 35 5. Add 50% of remaining water
6. Mix at low speed for 0.5 minute, scrape down and mix at medium speed for 2 minutes
7. Add remaining water, mix at low speed for 0.5 minute, scrape down and mix at medium speed for 2 minutes
- 40 8. Scale batter into each of two greased pans
9. Bake at 175-190°C

Evaluation of Cakes

Cakes should be graded for volume and texture on the same day as baked according to AACC Method 10-90.

- 5 The internal structure may be scored for the uniformity and size of cells as well as thickness of the walls; the grain; texture, such as moisture, tenderness and softness; crumb colour; and flavour.

10 Preparation of Cookies

Cookies may be prepared according to AACC Method 10-50D (in Approved Methods of the American Association of Cereal Chemists, Ninth Edition, March 1995; AACC, St. Paul MN, USA).

15 Basic recipe

	Flour	225 g
	Water	16 g
	Dextrose Solution	33 g
	Bicarbonate of Soda	2.5 g
20	Salt	2.1 g
	Sugar	130 g
	Shortening	64 g

Procedure

- 25 1. Cream shortening, sugar, salt and soda on low speed 3 minutes using an electric mixer (e.g., Hobart C-100).
2. Add dextrose solution and distilled water
3. Mix at low speed for 1 minute
4. Mix at medium speed for 1 minute
- 30 5. Add all flour and mix at low speed for 2 minutes
6. Scrape dough from bowl and place six portions at well-spaced points on lightly greased cookie sheet
7. Flatten dough lightly
8. Cut dough with cookie cutter
- 35 9. Bake at 205°C for 10 minutes

Evaluation of Cookies

Cookie width should be measured after cooling 30 minutes and can be done by the method according to AACC Method 10-50D.

- 40 The width of each of the six cookies is measured in mm, then

rotated 90° and remeasured to obtain the average width (W). An average thickness (T) may be obtained by measuring the cookies stacked on top of one another, then restacked in a different order. The spread factor is the ratio of W/T. However, the most sensitive and reliable estimate is the width measurement, and in some cases, thickness. Because the spread factor is a ratio of 2 empirically determined parameters, different values of W and T can result in the same W/T.

10 Preparation of Biscuits

Biscuits may be prepared according to AACC Method 10-31B (in Approved Methods of the American Association of Cereal Chemists, Ninth Edition, March 1995; AACC, St. Paul MN, USA).

15 Basic recipe

Flour	228 g
Shortening	40 g
Milk Solution ¹	135 g
Bicarbonate of Soda ²	3.4 g
20 Salt ²	4.5 g
Monocalcium Phosphate ²	130 g

¹50 g milk powder in 450 ml water

²omit if self-rising flour is used; use 240 g of self-rising

25 flour

Procedure

1. Sift together flour and other dry ingredients (bicarbonate of soda, salt and monocalcium phosphate, if used)
2. Add shortening to flour mixture
- 30 3. Mix, using electric mixer (e.g., Hobart, Kitchen Aid or equivalent) with timer control, at speed 1 for 15 seconds
4. Mix at speed 1 for 3 minutes
5. Add milk solution and mix at speed 1 for 15 seconds
6. Roll out dough using floured rolling pin
- 35 7. Cut dough with floured cutter
8. Place 8 dough pieces 4 cm apart on ungreased baking sheet.
9. Bake at 232°C for 10 minutes

Evaluation of Biscuits

- 40 Upon removal from oven, biscuits should be removed from the

baking sheet and cooled for 30 minutes. Measurements of the eight biscuits can be made according to AACC Method 10-31B to obtain a total weight, a total diameter and a height at the top centre of each biscuit.

5

Preparation of Pie Shells

Pie shells may be prepared according to AACC Method 10-60 (in Approved Methods of the American Association of Cereal Chemists, Ninth Edition, March 1995; AACC, St. Paul MN, USA).

10

Basic recipe

	Flour	100%
	Shortening	60%
	Salt	3.5%
15	Water	30-64%

All percentages are by weight relative to the wheat flour, and all ingredients are at 10°C before mixing.

20 Dough Preparation

1. Sift flour twice
2. Add shortening to flour and cut for 5 minutes using electric mixer (e.g., Hobart, Kitchen Aid or equivalent) with timer control, on low speed
- 25 3. Dissolve salt in a portion of water
4. Add salt solution to flour-shortening mixture, together with additional water if necessary
5. Mix at low speed for 2 minutes
6. Store dough at 10°C for 24 hours

30

Empty shells

1. Scale, press dough into ball
2. Roll dough, fold and roll again
3. Fold and roll a third time
- 35 4. Lay dough sheet over an inverted pie tin
5. Trim dough and prick with fork
6. Let dry for 30 minutes and cover with a second pan pressed down firmly
7. Bake at 218°C for 20-25 minutes, removing second pan after 10
- 40 minutes in the oven

Filled pies

1. Scale and roll bottom crust as outlined above for empty pie shell
- 5 2. Press dough sheet into pie tin and fill with either artificial fruit acid filling (water, corn starch, sugar and citric acid crystals) or true fruit filling (cling peaches, sugar corn starch and water)
3. Scale and roll dough once for top crust
- 10 4. Place over filling, trim and cut centre lightly
5. Press edge over wetted edge of bottom crust
6. Bake at 218°C for about 30 minutes.

Evaluation of Pie Crusts

- 15 Viscosity may be evaluated according to AACC Method 56-80. Other parameters of empty and filled pie shells may be measured according to AACC Method 10-60 24 hours and 12 or 16 hours after baking, respectively. Pie crusts may be evaluated empirically for whether they are baked through; the edges have
- 20 shrunk from edge of pan; blisters have appeared; the texture is flaky; the mouth-feel is tender; whether they are crisp or soft; the colour; and if the fruit filling has penetrated the crust.

Testing of Doughs and Breads

According to the methods of the present invention, the effect of adding an XET may be tested in doughs and breads by using the following method:

Recipe:

30	Water	60%
	Wheat Flour	100%
	Yeast	4%
	Salt	1.5%
	Sugar	1.5%

35

Procedure for Preparation of Breads

1. Dough mixing (Spiral mixer)

3 minutes at low speed

8 minutes at high speed

The mixing time may be adjusted by the skilled baker to obtain an optimum dough consistency under the testing conditions used.

2. 1st proof: 30°C - 80% RH, 20 minutes

3. Scaling and shaping;

4. Final proof: 32°C - 80% RH, 40 minutes;

5. Baking: 225°C, 20 minutes for rolls and 30 minutes for loaf.

Evaluation of Dough and Baked Products

Dough and baked products made from the straight dough method described above may be evaluated as follows for loaf specific volume, dough stickiness, dough firmness, dough extensibility, dough elasticity, crumb structure, and gluten strength.

Loaf specific volume: The mean value of 4 loaves volume is measured using the traditional rapeseed displacement method. The specific volume is calculated as volume per weight of bread (ml/g). The specific volume of the control (without enzyme) is defined as 100. The relative specific volume index is calculated as follows:

specific vol. of 4 loaves

Specific vol. index = ----- x100

spec. vol. of 4 control loaves

The dough stickiness, firmness, extensibility, elasticity and crumb structure may be evaluated relative to controls by the skilled test baker according to the following scales:

	1	2	3	4	5	6
Dough Stickiness	almost liquid	too sticky	sticky	normal	dry	not used
Crumb Structure	very poor	poor	non-uniform	uniform/good	very good	not used
Dough Firmness	extremely soft	too soft	soft/good	normal	firm	too firm
Dough Extensibility	too short	short	normal	good	long	too long

Dough stability / Shock test: After the second proof a pan containing the dough is dropped from a height of 20 cm. The dough is baked and the volume of the resulting bread is determined.

5

Gluten Strengthening: The strengthening effect of a given dough conditioner on wheat flour dough or gluten dough may be measured by dynamic rheological measurements. These measurements are able to show the strength of a dough under oscillation. Both wheat flour dough and gluten dough are viscoelastic materials. In oscillatory measurements, the viscoelastic properties of a wheat dough and a gluten dough can be divided into two components, the dynamic shear storage modulus G' and the dynamic shear loss modulus G'' . The ratio of the loss and the storage moduli is numerically equal to the tangent of the viscoelastic phase angle δ (Delta). An increase in the storage modulus G' and a decrease in the phase angle δ indicate a stronger and more elastic dough.

20 Example 1: Cloning and Expression of an XET from *Tiarosporella phaseolina* CBS No. 446.97

Chemicals used as buffers and substrates were commercial products of at least reagent grade.

25

Deposited organism:

Tiarosporella phaseolina CBS No. 446.97

Other strains:

- 30 W3124: a *Saccharomyces cerevisiae* strain (MATa; ura 3-52; leu 2-3, 112; his 3-D200; pep 4-1137; prcl::HIS3; prbl::LEU2; cir+) as described by van den Hazel, H.B; Kielland-Brandt, M.C.; Winther, J.R. Eur. J. Biochem. 207: 277-283, 1992.
- 35 DH10B: an *E. coli* strain (Life Technologies, Rockville MD, USA)

Plasmids:

pHD414: an *Aspergillus* expression vector derived from the plasmid p775 (EP 238 023) and described in WO 93/11249

pYES 2.0: (Invitrogen, San Diego CA, USA)

- 5 p3SR2: an *Aspergillus niger* plasmid containing the amdS gene (Christiensen, T., et al., 1988. Bio/Technology 6: 1419-22)

Media:

- 10 YPD: 10 g yeast extract, 20 g peptone, H₂O to 900 ml; autoclaved; 100 ml 20% glucose (sterile filtered) added

YPM: 10 g yeast extract, 20 g peptone, H₂O to 900 ml; autoclaved; 100 ml 20% maltodextrin (sterile filtered) added

- 15 SC-URA: 100 ml 10x Basal salt (75 g yeast nitrogen base, 113 g succinic acid, 68 g NaOH, H₂O ad 1000 ml, sterile filtered), 28 ml 20% casamino acids without vitamins, 10 ml 1% tryptophan, H₂O ad 900 ml, autoclaved, 3.6 ml 5% threonine and 100 ml 20% glucose or 20% galactose added

20 SC-agar: SC-URA; 20 g/l agar added

Molecular Biology Methods

- Expression cloning in yeast, extraction of RNA, cDNA synthesis, mung bean nuclease treatment, blunt-end formation with T4 DNA polymerase, and construction of libraries were performed as described by H. Dalboege et al. (1994, Mol. Gen. Genet. 243: 253-260.; WO 93/11249; WO 94/14953, which are hereby incorporated by reference).

30

Assay for XET Activity

- The assay uses an "XET-paper," a xyloglucan-coated paper which is dipped in a solution of labelled oligosaccharide, which is described in WO 97/11193. The procedure includes a method for preparing a labelled oligosaccharide and preparing xyloglucan-coated paper.
- 35

Preparation of labelled oligosaccharide

One gram of the reducing oligosaccharide 4-Q-[4-Q-[4-Q-[6-Q-
α-D-xylopyranosyl-β-D-glucopyranosyl]-6-Q-(2-Q-β-D-
galactopyranosyl)-α-D-xylopyranosyl-β-D-glucopyranosyl]-6-Q-(2-
5 Q-β-D-galactopyranosyl)-α-D-xylopyranosyl-β-D-glucopyranosyl]-D-
glucose ("XLLG") is dissolved in 25 ml of a saturated aqueous
solution of ammonium hydrogencarbonate containing 1 gram of
sodium cyanoborohydride (NaCNBH₃) and incubated in the dark at
25°C for 7 days to permit reductive amination. The ammonium
10 hydrogencarbonate is then removed by evaporation, and the
(ninhydrin-reactive) aminated derivative of XLLG is purified,
e.g., by gel-permeation chromatography or cation-exchange
chromatography. The product is believed to be an
oligosaccharidyl-1-amino-1-deoxyalditol, i.e., a derivative of
15 XLLG in which the reducing terminal D-glucose moiety has been
replaced by 1-amino-1-deoxy- D -glucitol.

Fifty mg of oligosaccharidyl-1-amino-1-deoxyalditol is
dissolved in 3 ml of 3% borax (disodium tetraborate; pH 9.0-9.5)
and a freshly prepared solution of 10 mg lissamine rhodamine
20 sulphonyl chloride (LSRSC) (Molecular Probes Inc., Eugene OR,
USA) in 0.75 ml of dry dimethylformamide (DMF) is added
gradually, with stirring, and the mixture is incubated in the
dark overnight. An additional 0.75 ml of LSRSC-DMF solution is
added and the mixture incubated for eight hours. The bright pink
25 oligosaccharidyl-1-amino-1-deoxyalditol-lissamine-rhodamine
conjugate (XLLGol-SR) is purified by gel-permeation
chromatography followed by reversed-phase chromatography on a
C₁₈-silica gel column. After the column is washed with water, a
methanol gradient is applied; the XLLGol-SR elutes in about 50%
30 methanol.

Preparation of xyloglucan-impregnated paper

Whatman No.1 filter paper is moistened with a 1% aqueous
solution of xyloglucan and dried. The XLLGol-SR preparation is
35 diluted in 75% aqueous acetone to give an absorbance at 580 nm
(A₅₈₀) of 0.2. The xyloglucan-coated paper is then dipped in the
XLLGol-SR solution and re-dried, resulting in "XET-paper".

Assay Procedure

An aliquot of the solution to be tested for XET activity is spotted onto XET-paper. Before the test sample dries, the sheet
5 is quickly sandwiched between two sheets of plastic (e.g., clear acetate sheets) and incubated, e.g., at 20°C for 1 hour. The incubated XET-paper with the plastic backing is then placed (paper-side down) in a dish containing about 150 ml of a solvent, e.g., freshly prepared ethanol/formic acid/water (1:1:1
10 by volume) which removes from the paper any unreacted XLLGol-SR but not XLLGol-SR which has become incorporated into the xyloglucan. The XET-paper is then rinsed in running water for 5 minutes, then washed in approximately 100 ml of acetone for 5 minutes, and dried thoroughly. The paper is then examined under
15 a short-wavelength ultraviolet lamp (e.g., emitting at 254 nm). The presence of XET is seen as a pink (orange-fluorescing) spot, which can be quantified, e.g., by use of a scanning spectrofluorometer.

Culture Cultivation

Tiarosporella phaseolina CBS No. 446.97 was inoculated from a plate with outgrown mycelium into a shake flask containing 100 ml medium B (20 g soyabean meal, 5 g maltodex 01, (Roquette 101-7845), 15 g wheat bran, 3 g peptone (Difco 0118), 10 g
25 cellulose avicel (Merck 2331), 0.2 ml pluronic (PE-6100, 101-3068), 1 g olive oil, deionized water to 1000 ml). The culture was incubated at 26°C and 200 rpm for 7 days. The resulting culture broth was filtered through miracloth and the mycelia were stored in liquid nitrogen.

30

RNA Isolation

Total RNA was extracted from frozen mycelia using guanidinium thiocyanate followed by ultracentrifugation through a 5.7 M CsCl cushion. Poly(A)⁺RNA was purified by oligo(dT)-
35 cellulose affinity chromatography using the procedures described in WO 94/14953.

cDNA synthesis

Double-stranded cDNA was synthesised from 5 mg poly(A)⁺ RNA by the RNase H method (Gubler and Hoffman, 1983. Gene 25:263-269; Sambrook, J. et al., *supra*). The poly(A)⁺ RNA (5 µg in 5 µl of DEPC-treated water) was heated at 70°C for 8 min. in a pre-siliconized, RNase-free microfuge tube, quenched in ice, then 45 µl of reverse transcriptase buffer (50 mM Tris-Cl, pH 8.3, 75 mM KCl, 3 mM MgCl₂, 10 mM DTT, Bethesda Research Laboratories) containing 1 mM dATP, dGTP and dTTP and 0.5 mM 5-methyl-dCTP (Pharmacia Biotech, Uppsala SE), 40 units human placental ribonuclease inhibitor (RNasin, Promega Corp., Madison WI, USA), 1.45 µg of oligo(dT)₁₈-Not I primer (Pharmacia Biotech) and 1000 units SuperScript II RNase H. reverse transcriptase (Bethesda Research Laboratories) was added. First-strand cDNA was synthesised by incubating the reaction mixture at 45°C for 1 hour. After synthesis, the mRNA:cDNA hybrid mixture was filtered through a MicroSpin S-400 HR (Pharmacia Biotech) spin column according to the manufacturer's instructions.

The recovered hybrids were diluted in 250 µl second strand buffer (20 mM Tris-Cl, pH 7.4, 90 mM KCl, 4.6 mM MgCl₂, 10 mM (NH₄)₂SO₄, 0.16 mM βNAD⁺) containing 200 µl of each dNTP, 60 units *E. coli* DNA polymerase I (Pharmacia Biotech), 5.25 units RNase H (Promega Corp.) and 15 units *E. coli* DNA ligase (Boehringer Mannheim GmbH, Mannheim DE). Second strand cDNA synthesis was achieved by incubating the reaction tube at 16°C for 2 hours and an additional 15 min. at 25°C. The reaction was stopped by addition of EDTA to a final concentration of 20 mM followed by phenol and chloroform extractions.

Mung bean nuclease treatment

The double-stranded cDNA was precipitated at -20°C for 12 hours by addition of 2 vols 96% EtOH, 0.2 vol 10 M NH₄Ac, recovered by centrifugation, washed in 70% EtOH, dried and resuspended in 30 µl mung bean nuclease buffer (30 mM NaAc, pH 4.6, 300 mM NaCl, 1 mM ZnSO₄, 0.35 mM DTT, 2% glycerol) containing 25 units mung bean nuclease (Pharmacia Biotech). The

single-stranded hair-pin DNA was clipped by incubating the reaction at 30°C for 30 min., followed by addition of 70 µl 10 mM Tris-Cl, pH 7.5, 1 mM EDTA, phenol extraction and precipitation with 2 vols of 96% EtOH and 0.1 vol 3 M NaAc, pH 5.2 on ice for 30 min.

Blunt-end formation with T4 DNA polymerase

The double-stranded cDNAs were recovered by centrifugation and resuspended in 30 ml T4 DNA polymerase buffer (20 mM Tris-acetate, pH 7.9, 10 mM MgAc, 50 mM KAc, 1 mM DTT) containing 0.5 mM of each dNTP and 5 units T4 DNA polymerase (New England Biolabs, Beverly MA, USA). The reaction mixture was incubated at 16°C for 1 hour, and stopped by addition of EDTA to a final concentration of 20 mM, followed by phenol and chloroform extractions, and precipitation for 12 hours at -20°C by addition of 2 vols 96% EtOH and 0.1 vol 3 M NaAc pH 5.2.

Adaptor ligation. Not I digestion and size selection

After the fill-in reaction the cDNAs were recovered by centrifugation, washed in 70% EtOH and dried. The cDNA pellet was resuspended in 25 µl ligation buffer (30 mM Tris-Cl, pH 7.8, 10 mM MgCl₂, 10 mM DTT, 0.5 mM ATP) containing 2.5 µg non-palindromic BstXI adaptors (Invitrogen Corp., San Diego CA, USA) and 30 units T4 ligase (Promega Corp.) and incubated at 16°C for 12 hours. The reaction was stopped by heating at 65°C for 20 min. and then cooling on ice for 5 min. The adapted cDNA was digested with Not I (New England Biolabs) by addition of 20 µl water, 5 µl 10x Not I restriction enzyme buffer (New England Biolabs) and 50 units enzyme, followed by incubation for 2.5 hours at 37°C. The reaction was stopped by heating at 65°C for 10 min. The cDNAs were purified by gel electrophoresis on a 0.8% SeaPlaque GTG low melting temperature agarose gel (FMC Corp. Bioproducts, Rockland ME, USA) in 1x TBE. The cDNAs were size-selected at a cut-off of 0.7 kb and recovered from the gel by use of b-Agarase (New England Biolabs) according to the manufacturer's instructions and precipitated for 12 hours at -20°C by adding 2 vols 96% EtOH and 0.1 vol 3 M NaAc pH 5.2.

Construction of a cDNA Library

The directional, size-selected cDNAs were recovered by centrifugation, washed in 70% EtOH, dried and resuspended in 30 μ l 10 mM Tris-Cl, pH 7.5, 1 mM EDTA. The cDNAs were desalted through a MicroSpin S-300 HR (Pharmacia Biotech) spin column according to the manufacturer's instructions. Three test ligations were performed in 10 μ l ligation buffer (30 mM Tris-Cl, pH 7.8, 10 mM MgCl₂, 10 mM DTT, 0.5 mM ATP) containing 5 μ l double-stranded cDNA (reaction tubes #1 and #2), 15 units T4 ligase (Promega) and 30 ng (tube #1), 40 ng (tube #2) and 40 ng (tube #3, the vector background control) of BstXI-NotI cleaved pYES 2.0 vector. The ligation procedure was incubation at 16°C for 12 hours, followed by heating at 70°C for 20 min. and addition of 10 μ l water to each tube. One μ l of each ligation mixture was electroporated into 40 μ l electrocompetent *E. coli* DH10B cells (Life Technologies, Inc., Rockville MD, USA) as described by Sambrook, et al., *supra*.

Transformed *E. coli* were plated onto LB+ampicillin agar at a density resulting in 15,000-30,000 colonies/plate after incubation at 37°C for 24 hours. A cell suspension was obtained by adding 20 ml LB+ampicillin to the plate, then shaken in a 50 ml tube for 1 hour at 37°C. Plasmid DNA was isolated from the cells using the QIAGEN plasmid kit (Qiagen GmbH, Hilden DE) according to the manufacturer's instructions, then stored at -20°C.

One- μ l aliquots of purified plasmid DNA (100 ng/ml) from individual pools were transformed into *S. cerevisiae* W3124 by electroporation (Becker and Guarante, 1991. *Methods Enzymol.* 194: 182-187). The transformants were then plated on SC-agar containing 2% glucose and incubated at 30°C.

Identification of positive colonies

Colonies were screened indirectly for XET by finding xyloglucanase positive colonies. After 3-5 days of growth, the agar plates were replica plated onto SC-URA agar (with 20% galactose added) containing 0.1% AZCL xyloglucan ((Megazyme International Ireland Ltd., Bray County Wicklow, IE). The

plates were incubated for 3-7 days at 30°C. Xyloglucanase-positive colonies were identified as colonies surrounded by a blue halo. Cells from enzyme-positive colonies were plated onto agar for single colony isolation, and an enzyme-producing single colony was selected from each of the xyloglucanase-producing colonies identified. All xyloglucanase positive colonies were tested for XET and were found to be positive.

Expression of XET in *Aspergillus*

10 An XET-producing yeast colony was inoculated into 20 ml YPD broth in a 50 ml glass test tube, then incubated with shaking for 2 days at 30°C. The cells were harvested by centrifugation, and the DNA isolated as described in WO 94/14953. Transformation of *E. coli* and isolation of plasmid DNA were performed using standard procedures. The cDNA insert was excised using the restriction enzymes BamH I and Xba I and ligated into the *Aspergillus* expression vector pHD414, resulting in the plasmid pA2XG80.

20 The cDNA inset of pA2XG80 was sequenced using the Taq deoxy-terminal cycle sequencing kit (Perkin Elmer, Foster City CA, USA) and synthetic oligonucleotide primers on an ABI PRISM™ 377 DNA Sequencer (Applied Biosystems, Foster City CA, USA) according to the manufacturer's instructions.

25 The cDNA sequence encoding the XET gene is set forth in SEQ ID NO:1. The deduced amino acid sequence is set forth in SEQ ID NO:2.

Transformation of *Aspergillus oryzae*

30 *Aspergillus niger* protoplasts were prepared as described in WO 95/02043, p. 16, line 21 - page 17, line 12, which is hereby incorporated by reference. One hundred μ l of protoplast suspension was mixed with 5 - 25 μ g of the appropriate DNA in 10 μ l of STC (1.2 M sorbitol, 10 mM Tris-HCl, pH 7.5, 10 mM CaCl₂). Protoplasts were then mixed with p3SR2 (a plasmid containing a gene for *A. nidulans amdS*; Christensen, T., et al., 1988. Bio/Technology 6: 1419-1422) and incubated at ambient temperature for 25 minutes. 0.2 ml of 60% PEG 4000 (BDH

29576), 10 mM CaCl₂ and 10 mM Tris-HCl pH 7.5 was added and mixed, followed by adding and mixing in a second aliquot of 0.85 ml of the same solution, and a 25 minute incubation at room temperature. The mixture was centrifuged at 2500 g for 15 minutes and the pellet was resuspended in 2 ml of 1.2 M sorbitol. After a second sedimentation step the protoplasts were spread on minimal plates (Cove, 1966. Biochem. Biophys. Acta 113: 51-56) containing 1.0 M sucrose, pH 7.0, 10 mM acetamide and 20 mM CsCl. After incubation for 4 - 7 days at 37°C, spores were picked and spread for single colonies. This procedure was repeated to obtain spores of a single colony, which were then stored as defined transformants.

Positive transformants were confirmed by XET activity, assayed as described above, and by sequencing.

A clone, Cl.XG80, containing the XET cDNA sequence was deposited at Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ) on 24 February 1998 under Accession No. DSM 12032.

The invention described and claimed herein is not to be limited in scope by the specific embodiments herein disclosed, since these embodiments are intended as illustrations of several aspects of the invention. Any equivalent embodiments are intended to be within the scope of this invention. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

Various references are cited herein, the disclosures of which are incorporated by reference in their entireties.

CLAIMS

1. A method for preparing a dough and/or a baked product made from dough, comprising incorporating into the dough a xyloglucan endotransglycosylase (XET).
- 5 2. The method of claim 1, wherein the XET is obtained from a plant or microbial source.
3. The method of claim 2, wherein the XET is obtained from a strain of *Ascomycota*, preferably a strain of *Tiarosporella*.
4. The method of any of claims 1-3, wherein the XET is:
 - 10 (a) a polypeptide which is encoded by a nucleic acid sequence which hybridizes under low stringency conditions with (i) the nucleic acid sequence set forth in SEQ ID NO:1, (ii) its complementary strand, or (iii) a subsequence thereof;
 - (b) a polypeptide having an amino acid sequence which
15 has at least 50% identity with the amino acid sequence set forth in SEQ ID NO:2;
 - (c) an allelic form of (a) or (b);
 - (d) a fragment of (a), (b) or (c), wherein the fragment retains XET activity;
 - 20 (e) a polypeptide produced by a strain of *Tiarosporella phaseolina*; or
 - (f) a polypeptide encoded by the XET encoding part of the cDNA sequence cloned into DSM 12032.
- 25 5. The method of any of claims 1 to 4, wherein the XET is incorporated in an amount of about 0.01 mg to about 100 mg per kilogram of dough.
6. The method of claims 1 to 4, wherein the XET is incorporated in an amount which is effective for increasing
30 strength of the dough, increasing stability of the dough, reducing stickiness of the dough, improving machinability of

the dough, increasing volume of the baked product, improving crumb structure of the baked product, improving softness of the baked product, improving flavour of the baked product, or reducing staling of the baked product.

5 7. The method of any of claims 1 to 6, wherein the dough is obtained from one or more ingredients selected from the group consisting of wheat meal, wheat flour, corn meal, corn flour, durum flour, rye meal, rye flour, oat meal, oat flour, soy meal, soy flour, sorghum meal, sorghum flour, potato meal, and
10 potato flour.

8. The method of any of claims 1 to 7, wherein the dough is fresh or frozen.

9. The method of any of claims 1-8, wherein the baked product is selected from the group consisting of a bread, a
15 roll, a French baguette-type bread, a pasta, a noodle, a pizza, a pita bread, a tortilla, a taco, a cake, a pancake, a biscuit, a cookie, a pie crust, steamed bread, and a crisp bread.

10. The method of any of claims 1 to 9, further comprising incorporating one or more additional enzymes selected from the
20 group consisting of an amylase, a cellulase, a hemicellulase, a lipase, an oxidase, a pentosanase, a peptidase, a peroxidase, and a protease.

11. The method of claim 10, wherein the amylase is a *Bacillus stearothermophilus* maltogenic alpha-amylase.

25 12. The method of any of claims 1 to 11, further comprising incorporating one or more baking agents selected from the group consisting of a protein, an emulsifier, a granulated fat, an oxidant, an amino acid, a sugar, a salt, a flour, and a starch.

13. A dough which comprises a xyloglucan endotransglycosylase
30 (XET).

14. The dough of claim 13, wherein the effective amount of the XET is about 0.01 mg to about 100 mg per kilogram of dough.

15. A pre-mix for dough comprising flour and a xyloglucan endotransglycosylase (XET).

5 16. A baking additive in the form of a granulate or agglomerated powder, which comprises a xyloglucan endotransglycosylase (XET).

17. The baking additive of claim 16, wherein more than 95% by weight of the baking additive has a particle size between about
10 25 and about 500 mm.

SEQUENCE LISTING

<110> Novo Nordisk A/S

<120> Methods for Using Xyloglucan Endotransglycosylase in Baking

<130> 5235-WO, BLK

<140>

<141>

<160> 2

<170> PatentIn Ver. 2.1

<210> 1

<211> 975

<212> DNA

<213> Tiarosporella phaseolina

<220>

<221> CDS

<222> (45) .. (779)

<400> 1

ggatccgaat tccaactatc ctgccctcct ttcaagcgaa cacc atg aag ttc tcc 56

Met Lys Phe Ser

1

tcg gct ctg ttt ctg gcc gct acg gcg gtc ttg gct tcc gcc gcg ccg 104

Ser Ala Leu Phe Leu Ala Ala Thr Ala Val Leu Ala Ser Ala Ala Pro

5

10

15

20

ctt gag cgc cgc gcc gac ttt tgt ggt caa tgg gac aac gtg aag aac 152

Leu Glu Arg Arg Ala Asp Phe Cys Gly Gln Trp Asp Asn Val Lys Asn

25

30

35

gga cct tac act ctt tac aac aac ctg tgg gga aaa gat gct tcc gga 200

Gly Pro Tyr Thr Leu Tyr Asn Asn Leu Trp Gly Lys Asp Ala Ser Gly

40

45

50

gcc tcc gga tgc caa tgc acc ggc gtc gat agc ttc agc agc aac acc 248

Ala Ser Gly Ser Gln Cys Thr Gly Val Asp Ser Phe Ser Ser Asn Thr

55

60

65

atc gct tgg cac aca tcc tgg tcc tgg tcc ggt gct cag tac aat gtc 296

Ile Ala Trp His Thr Ser Trp Ser Trp Ser Gly Ala Gln Tyr Asn Val

70	75	80	
aag tct tac gca aac gtg gtc gtc gac atc acc tct aag aaa ctc agc			344
Lys Ser Tyr Ala Asn Val Val Val Asp Ile Thr Ser Lys Lys Leu Ser			
85	90	95	100
gcc atc agc agc att aac agc atc tgg cgc tgg gct tac acg ggt agc			392
Ala Ile Ser Ser Ile Asn Ser Ile Trp Arg Trp Ala Tyr Thr Gly Ser			
	105	110	115
aac att gtt gcc aat gtt gcc tac gat atc ttc act tca tcc act gtc			440
Asn Ile Val Ala Asn Val Ala Tyr Asp Ile Phe Thr Ser Ser Thr Val			
	120	125	130
ggg ggt agc gag gaa tat gaa atc atg ata tgg gtt ggt gct ctc ggt			488
Gly Gly Ser Glu Glu Tyr Glu Ile Met Ile Trp Val Gly Ala Leu Gly			
	135	140	145
ggg gct ggt ccg atc tca tct acc ggc tcc cct att gcc acc gtt tcc			536
Gly Ala Gly Pro Ile Ser Ser Thr Gly Ser Pro Ile Ala Thr Val Ser			
	150	155	160
ctt gca ggc tcc tcg tgg aag ctc tac aaa ggg ccc aac ggg cag atg			584
Leu Ala Gly Ser Ser Trp Lys Leu Tyr Lys Gly Pro Asn Gly Gln Met			
	165	170	180
acc gtg ttc agc ttc gtc gcc gag tcc aac gtg aac aac ttc agc ggt			632
Thr Val Phe Ser Phe Val Ala Glu Ser Asn Val Asn Asn Phe Ser Gly			
	185	190	195
gac ctt aac gct ttc atc aag tac ctc acc ggc aac cag ggc ctt ccc			680
Asp Leu Asn Ala Phe Ile Lys Tyr Leu Thr Gly Asn Gln Gly Leu Pro			
	200	205	210
gcc tcg caa tac atc aag agc att ggc gct ggc act gag ccg ttc acg			728
Ala Ser Gln Tyr Ile Lys Ser Ile Gly Ala Gly Thr Glu Pro Phe Thr			
	215	220	225
ggg tcc aac gcc aag ctg acc act tcc tcc tac act gtc agc ttc aga			776
Gly Ser Asn Ala Lys Leu Thr Thr Ser Ser Tyr Thr Val Ser Phe Arg			
	230	235	240
taa ctgtgaagct ttatgctgcc cttatgcac atccttgtac atagttatca			829
245			
ccaggggact cttgtaaata cgattgcctt attaacggcc tgcattctgct ttcacacaat			889

ggcatttacc aatcaacagt ggcctcgaa tccgtaaaag gtggcttaaa aaaaaaaaaa 949

aaaaaaaaaa aaattcctgc ggccgc

975

<210> 2

<211> 244

<212> PRT

<213> Tiarosporella phaseolina

<400> 2

Met Lys Phe Ser Ser Ala Leu Phe Leu Ala Ala Thr Ala Val Leu Ala

1 5 10 15

Ser Ala Ala Pro Leu Glu Arg Arg Ala Asp Phe Cys Gly Gln Trp Asp

20 25 30

Asn Val Lys Asn Gly Pro Tyr Thr Leu Tyr Asn Asn Leu Trp Gly Lys

35 40 45

Asp Ala Ser Gly Ala Ser Gly Ser Gln Cys Thr Gly Val Asp Ser Phe

50 55 60

Ser Ser Asn Thr Ile Ala Trp His Thr Ser Trp Ser Trp Ser Gly Ala

65 70 75 80

Gln Tyr Asn Val Lys Ser Tyr Ala Asn Val Val Val Asp Ile Thr Ser

85 90 95

Lys Lys Leu Ser Ala Ile Ser Ser Ile Asn Ser Ile Trp Arg Trp Ala

100 105 110

Tyr Thr Gly Ser Asn Ile Val Ala Asn Val Ala Tyr Asp Ile Phe Thr

115 120 125

Ser Ser Thr Val Gly Gly Ser Glu Glu Tyr Glu Ile Met Ile Trp Val

130 135 140

Gly Ala Leu Gly Gly Ala Gly Pro Ile Ser Ser Thr Gly Ser Pro Ile

145 150 155 160

Ala Thr Val Ser Leu Ala Gly Ser Ser Trp Lys Leu Tyr Lys Gly Pro

165 170 175

Asn Gly Gln Met Thr Val Phe Ser Phe Val Ala Glu Ser Asn Val Asn

180 185 190

Asn Phe Ser Gly Asp Leu Asn Ala Phe Ile Lys Tyr Leu Thr Gly Asn

195

200

205

Gln Gly Leu Pro Ala Ser Gln Tyr Ile Lys Ser Ile Gly Ala Gly Thr
210 215 220

Glu Pro Phe Thr Gly Ser Asn Ala Lys Leu Thr Thr Ser Ser Tyr Thr
225 230 235 240

Val Ser Phe Arg

PCT

5235-WO, SLK

Original (for SUBMISSION) - printed on 20.05.1999 11:10:39 AM

0-1	Form - PCT/RO/134 (EASY) Indications Relating to Deposited Microorganism(s) or Other Biological Material (PCT Rule 13bis)	
0-1-1	Prepared using	PCT-EASY Version 2.83 (updated 01.03.1999)
0-2	International Application No	
0-3	Applicant's or agent's file reference	5235-WO, SLK
1	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
1-1	page	8
1-2	line	35
1-3	Identification of Deposit	
1-3-1	Name of depositary institution	Centraalbureau voor Schimmelcultures
1-3-2	Address of depositary institution	Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, Netherlands
1-3-3	Date of deposit	28 January 1997 (28.01.1997)
1-3-4	Accession Number	CBS 446.97
1-4	Additional Indications	Tiarosporella phaseolina
1-5	Designated States for Which Indications are Made	all designated States
1-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE
2	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
2-1	page	9
2-2	line	35
2-3	Identification of Deposit	
2-3-1	Name of depositary institution	Centraalbureau voor Schimmelcultures
2-3-2	Address of depositary institution	Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, Netherlands
2-3-3	Date of deposit	28 January 1997 (28.01.1997)
2-3-4	Accession Number	CBS 448.97
2-4	Additional Indications	Zygomycota, Mucorales, Chaetocladiaceae
2-5	Designated States for Which Indications are Made	all designated States
2-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE
3	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
3-1	page	10
3-2	line	18

3-3	Identification of Deposit	
3-3-1	Name of depositary institution	Centraalbureau voor Schimmelcultures
3-3-2	Address of depositary institution	Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, Netherlands
3-3-3	Date of deposit	12 March 1996 (12.03.1996)
3-3-4	Accession Number	CBS 274.96
3-4	Additional Indications	Ascomycota, Loculoascomycetes, Dothidiales, Botrysphaeriacea
3-5	Designated States for Which Indications are Made	all designated States
3-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE
4	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
4-1	page	11
4-2	line	3
4-3	Identification of Deposit	
4-3-1	Name of depositary institution	Centraalbureau voor Schimmelcultures
4-3-2	Address of depositary institution	Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, Netherlands
4-3-3	Date of deposit	28 January 1997 (28.01.1997)
4-3-4	Accession Number	CBS 444.97
4-4	Additional Indications	Ascomycota, Discomycetes, Pezizales, Sarcosomataceae
4-5	Designated States for Which Indications are Made	all designated States
4-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE
5	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
5-1	page	12
5-2	line	4
5-3	Identification of Deposit	
5-3-1	Name of depositary institution	Centraalbureau voor Schimmelcultures
5-3-2	Address of depositary institution	Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, Netherlands
5-3-3	Date of deposit	28 January 1997 (28.01.1997)
5-3-4	Accession Number	CBS 447.97
5-4	Additional Indications	Ascomycota, Discomycetes, Leotiales, Sclerotiniaceae
5-5	Designated States for Which Indications are Made	all designated States
5-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE

PCT

5235-WO, SLK

Original (for SUBMISSION) - printed on 20.05.1999 11:10:39 AM

6	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
6-1	page	12
6-2	line	12
6-3	Identification of Deposit	
6-3-1	Name of depositary institution	Centraalbureau voor Schimmelcultures
6-3-2	Address of depositary institution	Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, Netherlands
6-3-3	Date of deposit	28 January 1997 (28.01.1997)
6-3-4	Accession Number	CBS 445.97
6-4	Additional Indications	Ascomycota, Pyrenomycetes, Xylariales, Amphisphaeriaceae
6-5	Designated States for Which Indications are Made	all designated States
6-6	Separate Furnishing of Indications	NONE
	These indications will be submitted to the International Bureau later	
7	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
7-1	page	12
7-2	line	32
7-3	Identification of Deposit	
7-3-1	Name of depositary institution	Centraalbureau voor Schimmelcultures
7-3-2	Address of depositary institution	Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, Netherlands
7-3-3	Date of deposit	02 January 1996 (02.01.1996)
7-3-4	Accession Number	CBS 830.95
7-4	Additional Indications	Ascomycota, Pyrenomycetes, Hypocreales, Hypocreaceae
7-5	Designated States for Which Indications are Made	all designated States
7-6	Separate Furnishing of Indications	NONE
	These indications will be submitted to the International Bureau later	
8	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
8-1	page	13
8-2	line	14
8-3	Identification of Deposit	
8-3-1	Name of depositary institution	Centraalbureau voor Schimmelcultures
8-3-2	Address of depositary institution	Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, Netherlands
8-3-3	Date of deposit	28 January 1997 (28.01.1997)
8-3-4	Accession Number	CBS 442.97
8-4	Additional Indications	Ascomycota, Pyrenomycetes, Halosphaeriales, Halosphaeriaceae

8-5	Designated States for Which Indications are Made	all designated States
8-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE
9	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
9-1	page	13
9-2	line	24
9-3	Identification of Deposit	
9-3-1	Name of depositary institution	Centraalbureau voor Schimmelcultures
9-3-2	Address of depositary institution	Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, Netherlands
9-3-3	Date of deposit	23 January 1997 (23.01.1997)
9-3-4	Accession Number	CBS 424.97
9-4	Additional Indications	Ascomycota, Pyrenomycetes, Diaporthales, Valsaceae
9-5	Designated States for Which Indications are Made	all designated States
9-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE
10	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
10-1	page	13
10-2	line	31
10-3	Identification of Deposit	
10-3-1	Name of depositary institution	Centraalbureau voor Schimmelcultures
10-3-2	Address of depositary institution	Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, Netherlands
10-3-3	Date of deposit	23 January 1997 (23.01.1997)
10-3-4	Accession Number	CBS 425.97
10-4	Additional Indications	Ascomycota, Pyrenomycetes, Diaporthales, Valsaceae
10-5	Designated States for Which Indications are Made	all designated States
10-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE
11	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
11-1	page	15
11-2	line	24

PCT

5235-WO, SLK

Original (for SUBMISSION) - printed on 20.05.1999 11:10:39 AM

11-3	Identification of Deposit	
11-3-1	Name of depositary institution	Centraalbureau voor Schimmelcultures
11-3-2	Address of depositary institution	Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, Netherlands
11-3-3	Date of deposit	28 January 1997 (28.01.1997)
11-3-4	Accession Number	CBS 443.97
11-4	Additional Indications	Basidiomycota, Hymenomycetes, Schizophyllales, Schizophyllac.
11-5	Designated States for Which Indications are Made	all designated States
11-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE
12	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
12-1	page	16
12-2	line	21
12-3	Identification of Deposit	
12-3-1	Name of depositary institution	DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH
12-3-2	Address of depositary institution	Mascheroder Weg 1b, D-38124 Braunschweig, Germany
12-3-3	Date of deposit	12 February 1997 (12.02.1997)
12-3-4	Accession Number	DSMZ 11404
12-4	Additional Indications	Bacillus alcalophilus
12-5	Designated States for Which Indications are Made	all designated States
12-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE

FOR RECEIVING OFFICE USE ONLY

0-4	This form was received with the International application: (yes or no)	YES
0-4-1	Authorized officer	Claus Jørgensen clark

FOR INTERNATIONAL BUREAU USE ONLY

0-5	This form was received by the International Bureau on:	
0-5-1	Authorized officer	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 99/00277

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: A21D 2/24

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: A21D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0562836 A1 (TAKARA SHUZO CO.LTD.), 29 Sept 1993 (29.09.93) --	1-17
A	WO 9513384 A1 (UNILEVER PLC ET AL), 18 May 1995 (18.05.95) --	1-17
A	WO 9723683 A1 (NOVO NORDISK A/S), 3 July 1997 (03.07.97) --	1-17



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

27 Sept 1999

Date of mailing of the international search report

29-09-1999

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Eva Johansson/Els
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 99/00277

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	File WPI, Derwent accession no. 95-380073, DAINIPPON PHARM CO LTD: "A protein-xylo-glucan complex - prepd. by combining protein with xylo-glucan using aminocarbonyl reaction;" & JP,A,7258292, 951009, DW9549 --	1-17
A	STN International, File CAPLUS, CAPLUS accession no. 1992:422231, Document no. 117:22231, Fry, Stephen C. et al: "Xyloglucan endotrans- glycosylase, a new wall-loosening enzyme activity from plants", BiochemJ (1992), 282(3) 821-8 --	1-17
P,A	WO 9838288 A1 (NOVO NORDISK A/S), 3 Sept 1998 (03.09.98) -- -----	1-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

30/08/99

International application No.

PCT/DK 99/00277

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0562836 A1	29/09/93	AU 667706 B AU 3540593 A CA 2092366 A JP 6086670 A JP 7079778 A US 5516694 A US 5840550 A	04/04/96 30/09/93 27/09/93 29/03/94 28/03/95 14/05/96 24/11/98
WO 9513384 A1	18/05/95	AU 698844 B AU 8112994 A CA 2176133 A CZ 9601361 A EP 0728208 A HU 74598 A HU 9601232 D JP 9511121 T PL 317046 A SK 57696 A	12/11/98 29/05/95 18/05/95 11/12/96 28/08/96 28/01/97 00/00/00 11/11/97 03/03/97 01/10/96
WO 9723683 A1	03/07/97	AU 1365897 A EP 0868559 A PL 327608 A	17/07/97 07/10/98 21/12/98
WO 9838288 A1	03/09/98	AU 6205198 A	18/09/98